

Sacramento Valley Groundwater Basin, Antelope Subbasin

- Groundwater Basin Number: 5-21.54
- County: Tehama
- Surface Area: 18,710 acres (29 square miles)

Basin Boundaries and Hydrology

The Antelope Subbasin comprises the portion of the Sacramento Valley Groundwater Basin bounded on the west by the Sacramento River, on the north by the Red Bluff Arch, on the northeast by the Cascade Range, and the southeast by Antelope Creek. The Antelope Subbasin is contiguous with the Dye Creek Subbasin to the south. Annual precipitation in the subbasin ranges from 23- to 27-inches, increasing to the east.

Hydrogeologic Information

Water-Bearing Formations

The aquifer system in this subbasin is comprised of continental deposits of Tertiary to late Quaternary age. The Quaternary deposits include Pleistocene Modesto and Riverbank Formations. The Tertiary deposits include the Pliocene Tehama Formation and the Tuscan Formation. The Tuscan Formation is the primary water producing zone in the basin.

Pleistocene Modesto Formation. The Pleistocene Modesto Formation (deposited between 14,000 to 42,000 years ago) consists of poorly indurated gravel and cobbles with sand, silt and clay derived from reworking and deposition of the Tehama, Tuscan, and Riverbank Formations. Well logs for wells drilled on the floodplain east of Red Bluff indicate that coarse grained clean sand and gravel extend to a depth of approximately 50 feet below the surface. Below this depth, cemented gravel, sandstone, and hard clay of the Tehama and Tuscan Formations are encountered (Omsted and Davis 1961). The Modesto Formation yields limited groundwater due to its limited thickness (DWR 1987).

Pleistocene Riverbank Formation. The Pleistocene Riverbank Formation (deposited between 130,000 and 450,000 years ago) is observed in the far northern extents of the subbasin. The Riverbank Formation yields limited groundwater due to its limited thickness and areal extents.

Pliocene Tuscan Formation. The Tuscan Formation is composed of volcanic breccia, tuff, tuff breccia, volcanic sandstone and conglomerate, basalt flows, and tuffaceous silt and clay. The formation is mostly consolidated tuff in the area of exposure east of the valley in the Cascade Range foothills. From there tuff breccias grade westerly into volcanic sands, gravels, and clay (DWR 1978). The Tuscan Formation is the major water-bearing aquifer in the northeastern portion of the Sacramento Valley. Thickness of the formation within the subbasin is approximately 1,500 feet (DWR 1987).

Pliocene Tehama Formation. The Tehama Formation interfingers with the Tuscan Formation along the Sacramento River and is exposed in westside

Sacramento River banks. The formation consists of fluvial deposits of predominantly silt and clay with gravel and sand interbeds (DWR 1987). The formation is identified within the subbasin at depths ranging from 100- to 150- feet (DWR 1987).

Recharge Areas

Recharge is from inflow from the Sacramento River, Salt Creek, and Antelope Creek. In an investigation conducted by U.S. Bureau of Reclamation, the upper and intermediate aquifer zones (located between the local groundwater elevation and 150 feet in depth) intercept the Sacramento River. Diurnal fluctuations in river stage produce diurnal water level fluctuations in the deeper aquifer zone (Ely 1994).

Restrictive Structures

The Inks Creek fold system is a series of northeast-trending folds north of the Antelope Subbasin. The system isolates the Redding Groundwater Basin from the Sacramento Valley Basin. The fold system is a hydrologic drainage divide and separates the Red Bluff Arch from the Chico Monocline (DWR 1987).

Groundwater Level Trends

Review of hydrographs for long-term comparison of spring-spring groundwater levels indicates a decline of 5- to 10-feet associated with the 1976-77 and 1987-94 droughts, followed by a recovery to pre-drought conditions of the early 1970's and 1980's. Generally, groundwater level data show a seasonal fluctuation of approximate 2- to 15-feet for normal and dry years. Overall, there does not appear to be any increasing or decreasing trends in groundwater levels.

Groundwater Storage

The storage capacity of the subbasin was estimated based on estimates of specific yield for the Sacramento Valley as developed in DWR (1978). Estimates of specific yield, determined on a regional basis, were used to obtain a weighted specific yield conforming to the subbasin boundary. The estimated specific yield for the subbasin is 7.2 percent. The estimated storage capacity to a depth of 200 feet is approximately 269,179 acre-feet.

Groundwater Budget (Type B)

Estimates of groundwater extraction for the Antelope Subbasin are based on a survey conducted by the California Department of Water Resources in 1994. The survey included landuse and sources of water. Estimates of groundwater extraction for agricultural and municipal/industrial uses are 17,000 and 2,100 acre-feet respectively. Deep percolation of applied water is estimated to be 3,800 acre-feet.

Groundwater Quality

Characterization. Groundwater in the subbasin is characterized as calcium-magnesium bicarbonate and magnesium-calcium bicarbonate. Total dissolved solids (TDS) range from 119- to 558- mg/L, averaging 280 mg/L (DWR unpublished data).

Impairments. High concentrations of boron, chloride, and TDS are found in groundwater in the vicinity of Salt Creek and Little Salt Creek. Nitrate concentrations of 20- to 45- mg/L have been observed within the west-central portion of the basin (DWR 1987).

Water Quality in Public Supply Wells

Constituent Group ¹	Number of wells sampled ²	Number of wells with a concentration above an MCL ³
Inorganics – Primary	17	0
Radiological	10	0
Nitrates	17	0
Pesticides	6	0
VOCs and SVOCs	3	0
Inorganics – Secondary	17	3

¹ A description of each member in the constituent groups and a generalized discussion of the relevance of these groups are included in *California's Groundwater – Bulletin 118* by DWR (2003).

² Represents distinct number of wells sampled as required under DHS Title 22 program from 1994 through 2000.

³ Each well reported with a concentration above an MCL was confirmed with a second detection above an MCL. This information is intended as an indicator of the types of activities that cause contamination in a given basin. It represents the water quality at the sample location. It does not indicate the water quality delivered to the consumer. More detailed drinking water quality information can be obtained from the local water purveyor and its annual Consumer Confidence Report.

Well Characteristics

Well yields (gal/min)		
Municipal/Irrigation	Range: 300 – 800	Average: 575 (4 Well Completion Report)
Total depths (ft)		
Domestic	Range: 40 - 450	Average: 104 (702 Well Completion Reports)
Municipal/Irrigation	Range: 40 - 600	Average: 176 (92 Well Completion Reports)

Active Monitoring Data

Agency	Parameter	Number of wells /measurement frequency
DWR	Groundwater levels	4 wells semi-annually
DWR	Miscellaneous water quality	5 wells biennially
Department of Health Services	Miscellaneous water quality	22

Basin Management

Groundwater management: Tehama County adopted a groundwater ordinance in 1994.
Tehama County adopted a countywide AB3030 plan in 1996.

Water agencies

Public Tehama County Flood Control and Water Conservation District, City of Red Bluff

Private

Selected References

California Department of Water Resources (DWR) in cooperation with the United States Geological Survey (USGS). 1978. Evaluation of Groundwater Resources: Sacramento Valley.

California Department of Water Resources. 1978. Evaluation of Groundwater Resources: Sacramento Valley. Department of Water Resources in cooperation with the United States Geological Survey. Appendix A. Bulletin 118-6.

California Department of Water Resources. April 1987. Antelope Groundwater Study. Northern District.

California Department of Water Resources. 2000. Geology and Hydrogeology of the Freshwater Bearing Aquifer Systems of the Northern Sacramento Valley, California. In Progress.

Helley EJ, Harwood DS. 1985. Geologic Map of the Late Cenozoic Deposits of the Sacramento Valley and Northern Sierran Foothills, California. USGS Map MF-1790.

Olmsted FH, Davis GH. 1961. Geologic Features and Ground Water Storage Capacity of the Sacramento Valley, California. USGS. Water Supply Paper 1497.

Bibliography

Bailey EH. 1966. Geology of Northern California. California Division of Mines and Geology. Bulletin 190.

Berkstressor CF. 1973. Base of Fresh Water in the Sacramento Valley and Sacramento-San Joaquin Delta, California. U.S. Geological Survey in Cooperation with California Department of Water Resources.

Bertoldi GT, Johnson RH, Evenson KD. 1991. Groundwater in the Central Valley, California - A Summary Report. Regional Aquifer System Analysis--Central Valley, California. USGS. Professional Paper 1401-A.

Beyer LA. 1993. Sacramento Basin Province. USGS.

Bryan K. 1923. Geology and Ground-water Resources of Sacramento Valley, California. USGS. 495.

California Department of Pesticide Regulation. 1993. Sampling for Pesticide Residues in California Well Water, 1993 Well Inventory Database. California Environmental Protection Agency.

California Department of Water Resources. 1958. Ground Water Conditions in Central and Northern California 1957-58. California Department of Water Resources. Bulletin 77-58.

California Department of Water Resources. 1964. Groundwater Conditions in Central and Northern California, 1961-62. California Department of Water Resources.

- California Department of Water Resources. 1964. Quality of Ground Water in California 1961-62, Part 1: Northern and Central California. California Department of Water Resources. Bulletin 66-62.
- California Department of Water Resources. 1966. Precipitation in the Central Valley. Coordinated Statewide Planning Program. California Department of Water Resources, Sacramento District. Office Report.
- California Department of Water Resources. 1975. California's Ground Water. California Department of Water Resources. Bulletin 118.
- California Department of Water Resources. 1975. Progress Report Sacramento And Redding Basins Groundwater Study. California Department of Water Resources, Northern and Central Districts, in cooperation with the U.S. Geological Survey. Bulletin 118.
- California Department of Water Resources. 1980. Ground Water Basins in California. California Department of Water Resources. Bulletin 118-80.
- California Department of Water Resources. 1987. Progress Report Sacramento and Redding Basins Ground Water Study. California Department of Water Resources, Northern and Central Districts, in cooperation with the U.S. Geological Survey.
- California Department of Water Resources. 1993. Ground Water Levels in the Sacramento Valley Ground Water Basin; Tehama County. California Department of Water Resources, Northern District.
- California Department of Water Resources. 1995. Sacramento Valley Groundwater Quality Investigation. California Department of Water Resources, Northern District.
- California Department of Water Resources. 1998. California Water Plan Update. California Department of Water Resources. Bulletin 160-98, Volumes 1 and 2.
- Cherven VB, Edmondson WF. 1992. Structural Geology of the Sacramento Basin: Annual Meeting, Pacific Section AAPG, Sacramento, California, April 27, 1992-May 2, 1992.
- Dickinson WR, Ingersoll RV, Graham SA. 1979. Paleogene Sediment Dispersal and Paleotectonics in Northern California. Geological Society of America Bulletin 90:1458-1528.
- Fogelman RP. 1976. Descriptions and Chemical Analysis for Selected Wells in the Central Sacramento Valley, California. USGS. OF-76-472.
- Fogelman RP. 1978. Chemical Quality of Ground Water in the Central Sacramento Valley, California. USGS. Water Resources Investigations 77-133.
- Fogelman RP. 1982. Dissolved-solids Concentrations of Groundwater in the Sacramento Valley, California. USGS. HA-645.
- Fogelman RP. 1983. Ground Water Quality in the Sacramento Valley, California, Water Types and Potential Nitrate and Boron Problem Areas. USGS. HA-651.
- Fogelman RP, Rockwell GL. 1977. Descriptions and Chemical Analysis for Selected Wells in the Eastern Sacramento Valley, California. USGS. OF-77-486.
- Harwood DS, Helley EJ. 1982. Preliminary Structure Contour Map of the Sacramento Valley, California, Showing Major Late Cenozoic Structural Features and Depth to Basement. USGS.
- Harwood DS, Helley EJ. 1987. Late Cenozoic Tectonism of the Sacramento Valley. USGS.
- Harwood DS, Helley EJ, Doukas MP. 1981. Geologic Map of the Chico Monocline and Northeastern Part of the Sacramento Valley, California. USGS.
- Hull LC. 1984. Geochemistry of Groundwater in the Sacramento Valley, California. Central Valley of California RASA Project. USGS. Professional Paper 1401-B.
- Lydon PA. 1969. Geology and Lahars of the Tuscan Formation, Northern California. The Geological Society of America.
- Mankinen EA. 1978. Paleomagnetic Evidence for a Late Cretaceous Deformation of the Great Valley Sequence, Sacramento Valley, California. USGS.

- Mitten HT. 1972. Estimated Ground-water Pumpage in the Northern Part of the Sacramento Valley, California, 1966-69. USGS.
- Mitten HT. 1973. Estimated Ground-water Pumpage in the Northern Part of the Sacramento Valley, California, 1970-71. USGS.
- Page RW. 1974. Base and Thickness of the Post-Eocene Continental Deposits in the Sacramento Valley, California. U.S. Geological Survey in cooperation with California Department of Water Resources. Water Resources Investigations 45-73.
- Page RW. 1986. Geology of the Fresh Groundwater Basin of the Central Valley, California, with Texture Maps and Sections. Regional Aquifer System Analysis. USGS. Professional Paper 1401-C.
- Planert M, Williams JS. 1995. Ground Water Atlas of the United States, Segment 1, California, Nevada. USGS. HA-730-B.
- Poland JF, Evenson RE. 1966. Hydrogeology and Land Subsidence, Great Central Valley, California, Geology of Northern California. California Division of Mines and Geology. 239-247 p.
- Steele WC. 1980. Quaternary Stream Terraces in the Northwestern Sacramento Valley, Glenn, Tehama, and Shasta Counties, California. USGS.
- Tehama County Flood Control and Water Conservation District. 1996. Coordinated AB 3030 Groundwater Management Plan. Tehama County Flood Control and Water Conservation District.
- Tehama County Flood Control and Water Conservation District. 1999. Coordinated AB 3030 Groundwater Management Plan, First Annual Report. Tehama County Flood Control and Water Conservation District.
- U.S. Geological Survey. 1981. Water Resources Data for California; Volume 4, Northern Central Valley Basins and the Great Basin from Honey Lake Basin to Oregon State Line. USGS.
- Williamson AK, Prudic DE, Swain LA. 1985. Groundwater Flow in the Central Valley, California. USGS. OF-85-345.
- Williamson AK, Prudic DE, Swain LA. 1989. Groundwater Flow in the Central Valley, California. Regional Aquifer-System Analysis--Central Valley, California. USGS. Professional Paper 1401-D.

Errata

Changes made to the basin description will be noted here.