

Sacramento Valley Groundwater Basin, Dye Creek Subbasin

- Groundwater Basin Number: 5-21.55
- County: Tehama
- Surface Area: 27,730 acres (43 square miles)

Basin Boundaries and Hydrology

The Dye Creek Subbasin comprises the portion of the Sacramento Valley Groundwater Basin bounded on the southwest by the Sacramento River, on the northwest by Antelope Creek, on the east by the Chico Monocline, and on the south by Mill Creek. The Chico Monocline is a geographic boundary with some recharge likely occurring east of the boundary line. The subbasin is contiguous with Antelope and Los Molinos subbasins at depth. Annual precipitation is approximately 17 inches.

Hydrogeologic Information

Water-Bearing Formations

The aquifer system is comprised of continental deposits of Tertiary to late Quaternary age. The Quaternary deposits include Holocene basin deposits and Pleistocene deposits of the Modesto and Riverbank formations and Pleistocene fanglomerate. The Tertiary deposits include Pliocene Tehama and Tuscan formations.

Holocene Basin Deposits. The basin deposits are exposed east of Highway 99, north and south of Dairyville, within the central portion of the subbasin. Basin deposits are the result of sediment-laden floodwaters rising above the natural levees of streams and rivers and spreading across low-lying areas. Thickness of the deposits has not been determined. The deposits generally have low permeability and yield low quantities of water to wells. The quality of groundwater produced from basin deposits is often poor (USBR 1960).

Pleistocene Modesto Formation. The Modesto Formation (deposited between 14,000 and 42,000 years ago) is observed along the western extents of the subbasin. The formation consists of undifferentiated terrace deposits of unconsolidated weathered and unweathered gravel, sand, silt and clay. Thickness of the unit can range from 0- to 150- feet (DWR 2000).

Pleistocene Riverbank Formation. The Riverbank Formation (deposited between 130,000 and 450,000 years ago) is exposed east of the Sacramento River north of Mill Creek. The formation is not a significant water-bearing formation due to its limited depth and areal extents.

Pleistocene Fanglomerate. The fanglomerate is observed along the eastern foothills and within the southern third of the subbasin. The formation is an alluvial fan deposit derived from erosion and deposition of volcanic mudflows of the Tuscan Formation and consists of poly lithic volcanic clasts set in weathered tuffaceous matrix. The fan deposits are poorly sorted and somewhat indurated to well cemented. Thickness of the fan deposits is up to

150 feet (Ely 1994). The fanglomerate is not sufficiently thick to produce large quantities of groundwater (Olmsted and Davis 1961).

Pliocene Tuscan Formation. The Tuscan Formation is composed of a series of volcanic breccia, tuff, tuff breccia, volcanic sandstone and conglomerate, basalt flows, and tuffaceous silt and clay layers. The formation is described as four separate but lithologically similar units, A through D (with Unit A being the oldest), which in some areas are separated by layers of thin tuff or ash units (Helley and Harwood 1985). Units A, B, and C are found within the subbasin and extend in the subsurface west to the Sacramento River. Surface exposures of Unit D appear along the east side of the subbasin and east of the subbasin boundary. The subsurface extents of Unit D is unknown.

Unit A is the oldest water bearing unit of the formation and is characterized by the presence of metamorphic clasts within interbedded lahars, volcanic conglomerate, volcanic sandstone and siltstone. Unit B is composed of fairly equal distribution of lahars, tuffaceous sandstone, and conglomerate. Unit C consists of massive mudflow or lahar deposits with some interbedded volcanic conglomerate and sandstone. In the subsurface, these low permeability lahars form thick, confining layers for groundwater contained in the more permeable sediments of Unit B. The Tuscan Formation reaches a thickness of 1,500 feet over older sedimentary deposits (DWR 2000). The slope of the formation averages approximately 2.5 degrees, east of the valley, and steepens sharply to 10 to 20 degrees southwestward towards the valley at the Chico Monocline (Olmsted and Davis 1961). The formation flattens beneath valley sediments.

Pliocene Tehama Formation. The Tehama Formation consists of fluvial deposits of predominantly silt and clay with gravel and sand interbeds and occurs in the subsurface along the western boundary of the subbasin (DWR 1987).

Groundwater Level Trends

Review of hydrographs for long-term comparison of spring-spring groundwater levels indicates a decline of 2- to 5-feet associated with the 1976-77 and 1987-94 droughts, followed by a recovery to pre-drought conditions of early 1970's and 1980's. Generally, groundwater level data show a seasonal fluctuation ranging from 2- to 10-feet for normal and dry years. Overall, there does not appear to be any increasing or decreasing trends in the groundwater levels.

Groundwater Storage

The storage capacity of the subbasin was estimated based on estimates of specific yield for the Sacramento Valley as developed in DWR (1978). Estimates of specific yield, determined on a regional basis, were used to obtain a weighted specific yield conforming to the subbasin boundary. The estimated specific yield for the subbasin is 6.0 percent. The estimated storage capacity to a depth of 200 feet is approximately 331,620 acre-feet.

Groundwater Budget (Type B)

Estimates of groundwater extraction for the Dye Creek Subbasin are based on a survey conducted by the California Department of Water Resources in 1994. The survey included landuse and sources of water. Estimates of groundwater extraction for agricultural and municipal/industrial uses are 9,300 and 680 acre-feet respectively. Deep percolation of applied water is estimated to be 3,200 acre-feet.

Groundwater Quality

Characterization. Groundwater in Antelope, Dye Creek, Los Molinos, and Vina subbasins is characterized as calcium-magnesium bicarbonate and magnesium-calcium bicarbonate. Total dissolved solids (TDS) range from 119- to 558-mg/L, averaging 280 mg/L (DWR unpublished data).

Water Quality in Public Supply Wells

Constituent Group¹	Number of wells sampled²	Number of wells with a concentration above an MCL³
Inorganics – Primary	2	0
Radiological	2	0
Nitrates	2	0
Pesticides	1	0
VOCs and SVOCs	0	0
Inorganics – Secondary	2	0

¹ A description of each member in the constituent groups and a generalized discussion of the relevance of these groups are included in *California's Groundwater – Bulletin 118* by DWR (2003).

² Represents distinct number of wells sampled as required under DHS Title 22 program from 1994 through 2000.

³ Each well reported with a concentration above an MCL was confirmed with a second detection above an MCL. This information is intended as an indicator of the types of activities that cause contamination in a given basin. It represents the water quality at the sample location. It does not indicate the water quality delivered to the consumer. More detailed drinking water quality information can be obtained from the local water purveyor and its annual Consumer Confidence Report.

Well Characteristics

	Well yields (gal/min)	
Municipal/Irrigation	Range: 200 –3300	Average: 890 (5 Well Completion Reports)
	Total depths (ft)	
Domestic	Range: 19 – 220	Average: 94 (432 Well Completion Reports)
Municipal/Irrigation	Range: 55 –597	Average: 188 (56 Well Completion Reports)

Active Monitoring Data

Agency	Parameter	Number of wells /measurement frequency
DWR	Groundwater levels	8 wells semi-annually
DWR	Miscellaneous water quality	1 well biennially
Department of Health Services	Miscellaneous water quality	3

Basin Management

Groundwater management: Tehama County adopted a groundwater management ordinance in 1994. Tehama County adopted a countywide AB3030 plan in 1996.

Water agencies

Public Tehama County Flood Control and Water Conservation District.

Private

Selected References

California Department of Water Resources. April 1987. Antelope Groundwater Study. Northern District.

California Department of Water Resources. 2000. Geology and Hydrogeology of the Freshwater Bearing Aquifer Systems of the Northern Sacramento Valley, California. In Progress.

California Department of Water Resources. 1978. Evaluation of Ground Water Resources: Sacramento Valley: Appendix A. Bulletin 118-6.

Ely KE. 1994. "An Evaluation of Aquifer Characteristics Within the Tuscan Formation, Northeastern Sacramento Valley, Tehama County, California." MS Thesis CSUC. (On file in Department of Water Resources, Groundwater Section, Red Bluff, CA.

Olmsted FH, Davis GH. 1961. Geologic Features and Ground Water Storage Capacity of the Sacramento Valley, California. USGS. Water Supply Paper 1497.

Bibliography

Bailey EH. 1966. Geology of Northern California. California Division of Mines and Geology. Bulletin 190.

Berkstressor CF. 1973. Base of Fresh Water in the Sacramento Valley and Sacramento-San Joaquin Delta, California. U.S. Geological Survey in Cooperation with California Department of Water Resources.

Bertoldi GT, Johnson RH, Evenson KD. 1991. Groundwater in the Central Valley, California - A Summary Report. Regional Aquifer System Analysis--Central Valley, California. USGS. Professional Paper 1401-A.

Beyer LA. 1993. Sacramento Basin Province. USGS.

Bryan K. 1923. Geology and Ground-water Resources of Sacramento Valley, California. USGS. 495.

California Department of Pesticide Regulation. 1993. Sampling for Pesticide Residues in California Well Water, 1993 Well Inventory Database. California Environmental Protection Agency.

- California Department of Water Resources. 1958. Ground Water Conditions in Central and Northern California 1957-58. California Department of Water Resources. Bulletin 77-58.
- California Department of Water Resources. 1964. Groundwater Conditions in Central and Northern California, 1961-62. California Department of Water Resources.
- California Department of Water Resources. 1964. Quality of Ground Water in California 1961-62, Part 1: Northern and Central California. California Department of Water Resources. Bulletin 66-62.
- California Department of Water Resources. 1966. Precipitation in the Central Valley. Coordinated Statewide Planning Program. California Department of Water Resources, Sacramento District. Office Report.
- California Department of Water Resources. 1975. California's Ground Water. California Department of Water Resources. Bulletin 118.
- California Department of Water Resources. 1975. Progress Report Sacramento And Redding Basins Groundwater Study. California Department of Water Resources, Northern and Central Districts, in cooperation with the U.S. Geological Survey. Bulletin 118.
- California Department of Water Resources. 1980. Ground Water Basins in California. California Department of Water Resources. Bulletin 118-80.
- California Department of Water Resources. 1987. Progress Report Sacramento and Redding Basins Ground Water Study. California Department of Water Resources, Northern and Central Districts, in cooperation with the U.S. Geological Survey.
- California Department of Water Resources. 1993. Ground Water Levels in the Sacramento Valley Ground Water Basin; Tehama County. California Department of Water Resources, Northern District.
- California Department of Water Resources. 1995. Sacramento Valley Groundwater Quality Investigation. California Department of Water Resources, Northern District.
- California Department of Water Resources. 1998. California Water Plan Update. California Department of Water Resources. Bulletin 160-98, Volumes 1 and 2.
- Cherven VB, Edmondson WF. 1992. Structural Geology of the Sacramento Basin: Annual Meeting, Pacific Section AAPG, Sacramento, California, April 27, 1992-May 2, 1992.
- Dickinson WR, Ingersoll RV, Graham SA. 1979. Paleogene Sediment Dispersal and Paleotectonics in Northern California. Geological Society of America Bulletin 90:1458-1528.
- Fogelman RP. 1976. Descriptions and Chemical Analysis for Selected Wells in the Central Sacramento Valley, California. USGS. OF-76-472.
- Fogelman RP. 1978. Chemical Quality of Ground Water in the Central Sacramento Valley, California. USGS. Water Resources Investigations 77-133.
- Fogelman RP. 1982. Dissolved-solids Concentrations of Groundwater in the Sacramento Valley, California. USGS. HA-645.
- Fogelman RP. 1983. Ground Water Quality in the Sacramento Valley, California, Water Types and Potential Nitrate and Boron Problem Areas. USGS. HA-651.
- Fogelman RP, Rockwell GL. 1977. Descriptions and Chemical Analysis for Selected Wells in the Eastern Sacramento Valley, California. USGS. OF-77-486.
- Fogleman RP. 1979. Chemical Quality of Ground Water in the Eastern Sacramento Valley, California. USGS.
- Harwood DS, Helley EJ. 1982. Preliminary Structure Contour Map of the Sacramento Valley, California, Showing Major Late Cenozoic Structural Features and Depth to Basement. USGS.
- Harwood DS, Helley EJ. 1987. Late Cenozoic Tectonism of the Sacramento Valley. USGS.
- Harwood DS, Helley EJ, Doukas MP. 1981. Geologic Map of the Chico Monocline and Northeastern Part of the Sacramento Valley, California. USGS.

- Helley EJ, Harwood DS. 1985. Geologic Map of the Late Cenozoic Deposits of the Sacramento Valley and Northern Sierra Foothills, California. USGS. Map MF-1790.
- Hull LC. 1984. Geochemistry of Groundwater in the Sacramento Valley, California. Central Valley of California RASA Project. USGS. Professional Paper 1401-B.
- Lydon PA. 1969. Geology and Lahars of the Tuscan Formation, Northern California. The Geological Society of America.
- Mankinen EA. 1978. Paleomagnetic Evidence for a Late Cretaceous Deformation of the Great Valley Sequence, Sacramento Valley, California. USGS.
- Mitten HT. 1972. Estimated Ground-water Pumpage in the Northern Part of the Sacramento Valley, California, 1966-69. USGS.
- Mitten HT. 1973. Estimated Ground-water Pumpage in the Northern Part of the Sacramento Valley, California, 1970-71. USGS.
- Page RW. 1974. Base and Thickness of the Post-Eocene Continental Deposits in the Sacramento Valley, California. U.S. Geological Survey in cooperation with California Department of Water Resources. Water Resources Investigations 45-73.
- Page RW. 1986. Geology of the Fresh Groundwater Basin of the Central Valley, California, with Texture Maps and Sections. Regional Aquifer System Analysis. USGS. Professional Paper 1401-C.
- Planert M, Williams JS. 1995. Ground Water Atlas of the United States, Segment 1, California, Nevada. USGS. HA-730-B.
- Poland JF, Evenson RE. 1966. Hydrogeology and Land Subsidence, Great Central Valley, California, Geology of Northern California. California Division of Mines and Geology. 239-247 p.
- Steele WC. 1980. Quaternary Stream Terraces in the Northwestern Sacramento Valley, Glenn, Tehama, and Shasta Counties, California. USGS.
- Tehama County Flood Control and Water Conservation District. 1996. Coordinated AB 3030 Groundwater Management Plan. Tehama County Flood Control and Water Conservation District.
- Tehama County Flood Control and Water Conservation District. 1999. Coordinated AB 3030 Groundwater Management Plan, First Annual Report. Tehama County Flood Control and Water Conservation District.
- U.S. Geological Survey. 1981. Water Resources Data for California; Volume 4, Northern Central Valley Basins and the Great Basin from Honey Lake Basin to Oregon State Line. USGS.
- Williamson AK, Prudic DE, Swain LA. 1985. Groundwater Flow in the Central Valley, California. USGS. OF-85-345.
- Williamson AK, Prudic DE, Swain LA. 1989. Groundwater Flow in the Central Valley, California. Regional Aquifer-System Analysis--Central Valley, California. USGS. Professional Paper 1401-D.

Errata

Changes made to the basin description will be noted here.